



## Prevalence of Ixodid Ticks in Small Ruminants in and Around Jimma District

Mujahid Jewaro, \*Kufa Mustefa

Shashemene city Agricultural office Ethiopia

### Article Info

#### Publication Issue :

July-August-2023

Volume 6, Issue 4

#### Page Number : 41-51

#### Article History

Received : 01 Aug 2023

Published : 29 Aug 2023

**ABSTRACT** - A study on small ruminants tick was conducted in and around Jimmatown, Southwest Ethiopia, from November 2017 to April 2018 with the objectives of determining the tick infestation prevalence and identifying the common genera of hard ticks in indigenous breeds of small ruminants (sheep and goats). The study was conducted using cross-sectional design to assess the adult ticks attached on study animals. A total of 350 animals (299 sheep and 51 goats) were examined. From the total examined animals, 118 of them were found to harbour different tick genera, giving an overall prevalence of 33.7% (95% CI: 26.7- 38.9). The prevalence of tick infestation in goats and sheep was found to be 35.1% and 23.1 %, respectively. The prevalence of tick infestation between two age groups of animals were statistically insignificant ( $P>0.05$ ). However the prevalence was higher in young (53.3%) than adult (29.2%). The prevalence found to be statistically insignificant within species of small ruminants ( $P>0.05$ ). In this study , six genera of ticks were identified, with the following abundance among tick infested animals: Hyalomma(36.4%), Amblyomma(34.7%), Boophilus(22%), Ixodes6.7%) , Haemaphysalis(4.2%), and Reipicephalus (1.6%).The Preferred attachment sites for most of tick genera identifiedwere internal part of legs, scrotum/udder and anogenital in decreasing order. In conclusion small ruminants ticks infestation is highly prevalent in the study area. Therefore, attention should be given to the control and prevention of ticks and further study should be done to assess the seasonal dynamicity of ixodid ticks and tick borne diseases of small ruminants in the study area.

**Keywords** : Genera, Goat, Hard tick, Jimma, Sheep, survey

**Introduction** – Ethiopia is endowed with a very large and diverse livestock resource including sheep and goats, the country do have 29.33 million sheep, and 29.11 million goats (CSA, 2016).These small ruminants play a

**Copyright © 2023 The Author(s):** This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

significant role in socio-economic life of the people of Ethiopia (Minjauwet al., 2003). Sheep and goats account for 40% of cash income and 19% of the household meat consumption (Zelalem and Fletcher 1993). Owing to their high fertility, short generation interval and adaptation even in harsh environments, sheep, and goats are considered as investments and insurance to provide income to purchase food during seasons of crop failure and to meet seasonal purchases such as improved seed, fertilizer, and medicine for rural households. Hides and skins accounts for 12-16% of the total value of exports in Ethiopia (Minjauwet al., 2003a).

The current utilization of hides and skins in Ethiopia is estimated to be 45% for cattle hide, 75% goatskin, and 97% sheep skin with expected off take of 33, 35, and 7% for sheep, goats, and cattle, respectively. However, in recent years, this rank has been relegated to fifth level mainly because of rejection and down grading inflicted on hides and skin defects mainly due to infestation by external parasites (Kassa, 2006). Among the ectoparasitic infestations, ticks remain one of the most economically important parasites of livestock's in tropical and subtropical countries (Jongejan And Uilenberg, 1994). Ticks rank second to insects as vectors of transmissible diseases in man and animals (Opara 2016 and Ezeh, 2011). (Bowman et al., 1996) estimated.

more than 80% of world livestock population was infested by ticks, which were known to transmit viral, bacterial and protozoan pathogens causing Tick Borne Diseases (TBD) such as hemorrhagic fever, anaplasmosis, theileriosis and babesiosis (Rajput et al., 2006). Production losses due to ticks and tick-borne diseases (TBD) around the globe were put at US\$ 13.9 to US\$ 18.7 billion annually (De Castronet al., 1997).

In Ethiopia there are a lot of reports on the status of hard tick in small ruminants in different part of the country. For example several species of ticks belonging to genus *Amblyomma*, *Boophilus*, *Rhipicephalus*, *Hyalomma* and *Haemaphysalis* have been reported. Among these tick genera, the main ticks affecting small ruminants reported in Ethiopia were *Amblyomma* (40%), *Boophilus* (21%), *Haemaphysalis* (0.5%), *Hyalomma* (1.5%), and *Rhipicephalus* (37%) in different part of Ethiopia (Jaferet al., 2017), (Asmareet al ., 2012), (Serste and Wossen 2007). There are currently no studies on the prevalence and genera of ticks commonly affecting sheep and goats production in and around Jimma town, despite the fact that Jimma area is endowed with favorable weather condition suitable for the proliferation of ticks, thus the objectives of the study were:

- To determine the prevalence of tick infestation in small ruminants in the study area
- To identify the genera of ixodide ticks affecting sheep and goats

## II. MATERIALS AND METHODS

### 2.1. Description of study area

The study was conducted from November 2017 to April 2018 at Jimma town of Oromia Regional State south-western Ethiopia and Jimma University College of agriculture and veterinary medicine (JUCAVM). Jimma town is located at 355 km south-western of Addis Ababa, lies between 36° 10' E longitudes and 7° 40' N latitude an elevation ranging from 880 to 3360 meters above sea level (JZMSR 2004). Very currently Jimma Zone is divided into 17 districts (hosting a total population of over 2.4 million, (CSA 2008). The area is characterized by a humid tropical climate of heavy annual rainfall that ranges from 1200-2000 mm per year. About 70% of the total annual rainfall is received during rainy season, which lasts from the end of May to early September. The mean annual maximum and minimum temperature ranges from 25°C-30°C and 7°C-12°C, (OPEDJZ, 2002).

## 2.2. Study population

The study population comprises of local breeds of sheep and goats managed under extensive grazing system of both sex (male and female) in different Kebeles of Jimma town

## 2.3. Study Design

The study design was cross-sectional study, which was to assess the adult tick present in the body of small ruminants in the area

## 2.4. Study Methodology

Presence and absence ticks in the body of examined was done on the field, then adult ticks collection was done for identification of tick genera infesting sheep and goats.

### 2.4.1 Tick collection and identification

Adult ticks were collected from whole body regions of small ruminants into universal sample bottle containing 70% ethanol (Okello-Onenet al., 1999a and Walker et al., 2003). The body regions used for collections were head, neck, brisket, belly, udder/ scrotum, Anogenital region, leg and tail (Kaiseret al., 1982). Ticks were removed from the host skin whilst retaining their good condition for identification using good quality steel forceps. The collected adult ticks from each body regions were kept separately for identification in separate sample bottles. Then ticks transported to Jimma University, school of veterinary medicine, Veterinary parasatology laboratory. And identified using stereomicroscope following, in the laboratory the ticks collected were examined by seterio microscope. The morphology of the ticks was studied. Identification of the different genera of the ticks was accomplished with the help of the anatomical and morphological characteristics as described by (Okello-Onenet al., 1999a) and (Walker et al., 2003). During the collection time species (ovine/caprine), sex, age (young and adult) and body condition (Poor, Medium and Good) of animals were recorded.

## 2.5. Sampling and sample size

Examined animals were selected by systematic random sampling technique for ticks collection and identification from eight half-body regions of sheep and goats: Head, neck, Brisket, Belly, Udder or Scrotum, Anogenital, Leg and Tail Since there was no study conducted so far in Jimma town to determine the prevalence of tick infestation of small ruminants, as well as to get the maximum sample size (384), we considered an estimated prevalence of 50%, with 95% confidence interval and 5% absolute level of precision were considered according to the formula given by (Thrusfield 2005).

$$N = (1.96)^2 P_{exp}(1 - P_{exp}) / d^2$$

Where: N=Sample size, P<sub>exp</sub>= Expected prevalence (50%) and d = desired level of precision (5%). However a total 350 (299 sheep and 51 goats) animals were examined. A greater number of sheep was sampled because of their predominance in the area

## 2.6. Data Management and Analysis

The information obtaining from clinical examination, laboratory test and observation was entered on spreadsheet of Microsoft Excel. The data was screened for errors and coded before subjected to statistical analysis. Descriptive statistics used to analyze the data and percentages and tables were used to describe the results. To test the various risk factors associated with tick infestation, Chi-square test was used by using the Statistical Package STATA 11. In all the analyses, confidence level was held at 95% and p < 0.05 was considered as significant.

### III. RESULTS

#### 3.1. Overall prevalence of ectoparasites

From 350 small ruminant examined, 118 were infested by one or more genera ticks with an overall prevalence of 33.7% (95% CI: 26.7- 38.9). The overall tick infestation prevalence recorded in the male and female small ruminants of both species was 34.2% and 33.1% respectively, with no statistical significance variation (Table). Higher the tick infestation rate was observed in young (53.3%) small ruminants than adult (29.2 %) small ruminants, however the difference was not statistical significance ( $P > 0.05$ )

The overall prevalence of tick among sheep and goats were 35.1% (105/299) and 23.5% (12/51), respectively and no statistical significance variation was observed (Table1).

Table 1: Overall small ruminant's prevalence of tick infestation by sex, age and species of animals examined

Variable	No. of examined	No positive	Prevalence (%)	95% CI	X <sup>2</sup>	P-value
<b>Sex</b>						
Male	184	63	34.2	27.3- 41.6	0.47	0.827
Female	166	55	33.1	26.1- 40.8		
<b>Age</b>						
Young	13	7	53.3	25.1- 80.7	2.44	0.118
Adult	337	111	32.9	27.9- 38.2		
<b>Species</b>						
Ovine	299	105	35.1	29.7- 40.8	2.77	0.09
Caprine	51	12	23.5	12.7- 37.5		

#### 4.2 Prevalence of tick genera in small ruminants (sheep and goats)

During the study period six genera of different types of ticks were identified which are Amblyomma, Hyalomma, Boophilus, Haemaphysalis, Ixodes and Rhipicephalus. The overall prevalence the six genera of ixodide ticks among the small ruminants is depicted in table 2: The highest infestation was caused by Hyalomma (12.3%) followed by Amblyomma (11.71%), Boophilus (7.43%), Ixodes (2.3%), Haemaphysalis (1.2%) and Rhipicephalus (0.57%).

Table 2: The overall prevalence of genera of ticks identified in small ruminants (n= 350)

Genera of ticks	Number of	Prevalence (%)	95% CI	Chi-square, p-value
-----------------	-----------	----------------	--------	---------------------


**positive animals**

<i>Amblyomma</i>	41	11.71	8.8- 15.5	
<i>Boophilus</i>	26	7.43	4.5- 10.7	
<i>Haemaphysalis</i>	5	1.2	0.5- 3.3	75.4, 0.000, Df* = 5
<i>Hyalomma</i>	43	12.3	9.1- 16.2	
<i>Ixodes</i>	8	2.3	0.9- 4.4	
<i>Reipicephalus</i>	2	0.57	0.06- 2.1	

- Df : Degree of freedom

**1) 4.3. The proportion of genera of tick infestation in infested animals**

Out of all infested animals (n=118), the proportion of infection percentage by the six genera of tick identified and mixed infestation with more than one tick species is depicted in figure 4, 14.5% of infested animals were found to have a mixed infestation with more than one tick genera (figure 4).



**Figure4:** Relative abundance of tick genera during the study period among infested smallruminants (n=118)

## 2) 4.4. Assessment of genera of tick attachment sites

Each species of tick consistently favored various body regions of small ruminants (sheep and goats), but the preferences were stronger in some genera than others. The observed proportion of tick genera attachment sites during this study was summarized and shown on table 3. The most preferred predilection sites of all *Amblyomma* genera identified during this study was legs followed by scrotum, udder, anogenital, tail and head in decreasing order of preference. *Boophilus* preferred attachment sites in decreasing order were legs and brisket. During this study periods identified *Hyalomma* species were found primarily attached to legs and secondarily on scrotum or udder. *Haemaphysalis* and *Reipicephalus* mainly attached on legs (Table 3).

**Table 3.** Summary of the attachment sites percentages of the most common tick genera among infested small ruminants in the study areas

Body regions						
Tick genera	Head	Brisket	Scrotum/ Including ear	Tail	Anogenital	Leg+interdigita l space
<i>Amblyomma</i> (n=41)	1(2.4%)	-	7(17.1%)	3(7.3%)	4(9.7%)	25 (60.1%)
<i>Boophilus</i> (n=26)	-	3(11.%)	-	3(11.5)	2 (7.6%)	18(69.2%)
<i>Haemaphysalis</i> ( n=5)	-	-	1 (20%)	-	1(20%)	3 (60%)
<i>Hyalomma</i> (n=43)	1(2.3%)	2(4.6%)	3(6.8%)	-	1(2.3%)	37(86%)
<i>Ixodes</i> ( n=8)	1(12.5%)	-	1(12.5%)	-	1(12.5%)	5(62.5%)
<i>Reipicephalus</i> ( n=2)	-	-	-	-	-	2 (100%)

## DISCUSSION

The overall prevalence of tick infestation (33.7%) reported in this study is relatively moderate when compared to other reports from other areas of Ethiopia, for example the current finding was in agreement with the previous overall tick infestation prevalence report of 31.4% in and around Bahirdar town (Tesfayeet al., 2012). However a higher overall prevalence small ruminant tick infestation were reported elsewhere in this country; for instance 79.7%(Kedir and Petrose, 2015), 76.5% (Fufaet al., 2012) and 72.4% (Abebeet al., 2017) were reported in and around Bedele town, Fafen zone in Ethiopia Somali region and in and around Dire Dawa town. Relatively lower overall small ruminant tick infestation, which is 20% was reported in and around Gonder town (Fentahunet al., 2012) when compared with our finding. The difference in the prevalence might be due to the geographical difference, breed difference of the study animals and season of the study period.

The overall prevalence recorded in sheep (35.1%) and goats (23.5%). this study was close to the study conducted in and around Bahir Dar (Asmareet al., 2012) recorded an over prevalence of 37% in sheep and 23.9% in goats, similarly an overall prevalence of 21.2% and 17.1% in sheep and goats were recorded in and around Gonder town (Fentahunet al., 2012). Additionally (Yacobet al., 2008) reported very similar overall tick infestation prevalence in sheep (31.7%) and goat (18.3%) in Wolayita Sodo,

On the contrary the overall tick infestation rate recorded in sheep and goats in this study was lower than the reports done by (Seyoumet al., 2015) in Miesso district, Western Hararghe, who recorded a prevalence of 87.5 % (goats) and 89.9 % (sheep), (Eyob and Matios 2014) recorded a higher prevalence of 97.58% (goats) and 69.86% (sheep) in Dhas district of Borana pastoral area, Southern Rangelands of Ethiopia. Additionally 66.12% (goats) and 80.30% (sheep) over all prevalence was recorded in Bedelle district, Oromia Region, Ethiopia (Fufaet al., 2012). Environmental variations and differences in the time of year when the study was conducted could also contribute to differences in the prevalence of tick infestation in various areas of the country as temperature and relative humidity are the major ecological determinants for the reproduction and growth of tick populations (Latif and Walker, 2004).

The overall prevalence of tick in different age and sex groups of examined small ruminant was show no statistical significant variations ( $P > 0.05$ ). Even if there was no statically significant variations in young and adult animals examined in this study, the recorded prevalence in young animals (53.3%) seems large when compared with adult animals (32.9%). in contrary to this report (Fentahunet al., 2012 and Abebeet al., 2017) observed a significantly higher prevalence ( $P < 0.05$ ) in adult animals in their study in and around Gonder and Dire Dawa towns respectively.

The overall prevalence between male and female was statistically insignificant ( $P > 0.05$ ), this finding is in agreement with the study conducted in three districts of Fafen Zone in Somali Regional State, Eastern Ethiopia (Kedir and Petros, 2015). On the contrary the current findings, (Abebeet al., 2017) observed a statistically significant high prevalence in female small ruminants in their study. This study revealed the presence of six genera of hard ticks namely: *Amblyomma*, *Hyalomma*, *Boophilus*, *Haemaphysalis*, *Ixodes* and *Rhipicephalus*. These all indicates that the bionomic situation of the study area is favorable for the successive perpetuation of the pathogens transmitted by ticks and for their subsequent transmission to susceptible host.

In the present study, *hyalomma* had the highest frequency among the collected tick species in small ruminant (sheep and goats), the finding is in agreement with the previous studies done in different areas of Ethiopia (Kedir and Petros; 2015; Abebeet al., 2017). The least frequent genera of tick during the study period was *Rhipicephalus* (0.57%), similar find was reported in Arsi Area (Hailegebrielet al., 2015). In general the finding of

different genera of ticks in and around Jimma indicates the epidemiological significance in transmission of tick borne diseases such as anaplasmosis, babesiosis and rickettsial diseases.

The most preferred predilection sites of all Amblyommatinae identified during this study was legs followed by scrotum or udder, anogenital, tail and head in decreasing order of preference. Boophilus preferred attachment sites in decreasing order were legs and brisket. During this study periods identified Hyalomma species were found primarily attached to legs and secondarily on scrotum or udder. Haemaphysalis and Reipicephalus mainly attached on legs. The result of this study showed that the predilection sites of ticks over the body of small ruminants was high on legs, specifically in the interdigital spaces between hoof, this impairs productivity and mobility of the small ruminants (Sheep and goats).

#### CONCLUSION AND RECOMMENDATIONS

From this study result it is possible to conclude that the overall prevalence of tick infestation in sheep and goats is relatively high. Sex, age and species of small ruminant were not principally determinants in the overall tick infestation prevalence in the study area. Among the tick genera identified in the study areas the most abundant were Hyalomma, Amblyomma and Boophilus. The preferred attachment sites for most of tick genera identified were internal part of legs, scrotum/udder and anogenital in decreasing order. Therefore the following recommendations were put forward

- Strategic tick control: Application of acaricides aimed at reduction of ticks population based on information about their seasonal activity.
- Integrated tick control: These encompass biological, chemical and ecological control methods, should be used
- Extension work: Educating animal owners on the problems of tick, and the different control methods, which can be available in their areas. Successes of ticks control generally associated with good extension work.
- Tick-borne diseases assessment: Importance, distribution, and seasonal prevalence of tick borne survey should be conducted.

#### REFERENCES

- [1]. Abebe, H., Ahmed, J., Wendemagegn, D., Tsehay, A., Silesh, S., & Abebe, H. (2017): Prevalence Of Tick Infestation On Small Ruminants In And Around Dire Dawa, Eastern Ethiopia Journal of Parasitology and Vector Biology, 9(4), pp.27- 33.
- [2]. Asmare, A, Kassahun, A. & Fentahun, T. (2012). Occurrence of small ruminant ectoparasites in and around Bahir Dar, northwest Ethiopia. Advances in Biological Research, 6(5), 170-176.
- [3]. Bowman, D. M. J. S. (1996): "Diversity patterns of woody species on a latitudinal transect from the monsoon tropics to desert in the Northern Territory, Australia." Australian Journal of Botany 44.5 571-580.
- [4]. Central Statistical Agency (CSA) (2016): Federal democratic republic of Ethiopia.
- [5]. Central statistical agency. Agricultural sample survey, Volume II, Report on livestock and livestock characteristics. Statistical bulletin 583, Addis Ababa, Ethiopia.
- [6]. CSA, 2008. (Central Statistics Authority) (2008) Central Statistics Authority of the Federal Democratic Republic of Ethiopia.
- [7]. De Castro, J.J., (1997): Sustainable tick borne diseases control in Western Ethiopia. J. S. Afr. Vet. Assoc., 71(4): 240-243.



- [8]. Eyob, E., & Matios, L. (2014). Preliminary survey on the distribution of Ixodid ticks in small ruminants of dhas district of borena pastoral area, Southern Rangelands of Ethiopia. *Advanced in Biological Research*, 5, 87–91.
- [9]. Fentahun, Tewodros, Fasil Woldemariam, Mersha Chanie, and Malede Berhan. (2012):
- [10]. Prevalence of ectoparasites on small ruminants in and around Gondar Town." *American-Eurasian Journal of Scientific Research* 7, no. 3 106-111.
- [11]. Fufa., Abunna., Tura and. J., and Regassa, A. (2012b) Status of tick's infestation in small ruminants of Bedelle district, Oromia Region, Ethiopia. *Global Veterinaria*. 8(5): 459-462.
- [12]. Hailegebriel, Bedada, Terefe Getachew, and Hailu Tolossa Yacob., (2015): "Current status of ectoparasites in sheep and management practices against the problem in ectoparasites controlled and uncontrolled areas of Arsi zone in Oromia region, Ethiopia." *Journal of Veterinary Science and Technology* 6. Special Issue.
- [13]. Jafer Ahmed, Danel Wendemagegn, Abraham Tsehay, Samson Silesh, and Henok Abebe. (2017). "PREVALENCE OF TICK INFESTATION ON SMALL RUMINANTS IN AND AROUND DIRE DAWA, EASTERN ETHIOPIA." *International Journal of Research - Granthaalayah*, 5(5), 326-336.
- [14]. Jongejan, F. and Uilenberg, G. (1994): Ticks and control methods. *Ectoparasites of animals and control methods. Rev. Sci. tech. Off. Int. Epiz.*, 13(4), pp. 1201-1226.
- [15]. Jongejan, F. and Uilenberg, G. (2004): The Global Importance of Ticks. *Parasitology*, 129, S3-S14.
- [16]. JZMSR (Jimma Zone Meteorology Station Report), (2004): Ten year's climate data. JZMS. Jimma, Ethiopia., pp: 36.
- [17]. Kaiser MN, Sutherst RW, Bourne AS (1982) Relationship between ticks and zebu cattle in southern Uganda. *Trop Anim Health Prod* 14:63–74
- [18]. Kaiser, M.N, 1987. Ethiopia report on tick taxonomy and biology, AG; DP/ETH/83/23 consultant report- FAO of the United Nations pp 92.
- [19]. Kassa, B. (2006): 'Cockle, mange and pox: Major threats to the leather industry in Ethiopia. Ethiopian leather industry: Perseverance towards value addition', Proceedings of the National Workshop, Addis Ababa, Ethiopia, December 14–15, 2006. Pp: 71–92.
- [20]. Kedir Mohammed and Petros Admasu (2015): Prevalence of Ixodid Ticks in Small Ruminants in Selected Districts of Fafen Zone, Eastern Ethiopia *European Journal of Applied Sciences* 7(2): 50-55.
- [21]. Minjauw, B., and McLeod, A. (2003a): Tick borne diseases and poverty. The impact of ticks and tick borne diseases on livestock owners in India and Eastern health program center for tropical veterinary medicine, University of Edinburgh, UK. Pp: 24-57.
- [22]. Okello-Onen, J.; Hassan, S.M. and Essuman, S. (1999a): Taxonomy of African ticks, an identification manual. International Center for Insect Physiology and Ecology press, Nairobi, Kenya, pp.1-124.
- [23]. Opara, M. N. O., Opara, M. N., Nwachukwu, M., & Ezeh, N. O. (2016). Ixodid ticks of cattle in Borno and Yobe states of northeastern Nigeria: breed and coat colour preference. *Animal Research International*, 8(1).
- [24]. OPEDJZ (Office of Planning and Economic Development for Jimma Zone), (2002): Statistical Abstract. Jimma, Oromia, Ethiopi
- [25]. Sertse, T. and A. Wossene, (2007): A study on ectoparasites of sheep and goats in eastern part of Amhara region, Northeast Ethiopia. *Small Ruminant Research*, 69: 62-67.
- [26]. Seyoum, Z., Tadesse, T., & Addisu, A. (2015): Ectoparasites prevalence in small ruminants in and around Sekela, Amhara Regional State, Northwest Ethiopia. *Journal of veterinary medicine*, 2015.

- [27]. Tesfaye, D., Assefa, M., Demissie, T., &Taye, M. (2012): Ectoparasites of small ruminants presented at Bahir Dar Veterinary Clinic, Northwest Ethiopia. African Journal of Agricultural Research, 7(33), 4669-4674.
- [28]. Thrusfield, M. (2005): Veterinary Epidemiology, Second Edition. U.K: Blackwell .Science Ltd. W.B. Sounder Company. Pp: 1-78.
- [29]. Walker A, Bouattour A, Camicas J, Estrada-Pena A, Horak I, Latif A, (2003): Ticks of Domestic Animals in Africa: . Bioscience Report, Edinburgh. 1-221
- [30]. Yacob, H.T., A.T. Yalow and A.A. Dink, (2008): Ectoparasites prevalence in sheep and in goats in and around WolaitaSoddo, Southern Ethiopia. American-Eurasian Journal of Agriculture and Environmental Scientific Research, 159: 450-454.
- [31]. Zelalem, A., and Fletcher, I., C., (1993): Small ruminant productivity in the central Ethiopian mixed farming system. In proceedings of the 4th National Livestock Improvement conference, 13–15 November IAR, AddisAbaba, Ethiopia.

## Annexes

### Annex 1 : Sample collecting format

Animal no	Animal Ssp	Breed	Sex	Age	Site of adult tick	Adult tick	TickSpp	Lab. Result

### Annex 2: Estimated age for sheep and goats with different numbers of erupted permanent incisors

No. of permanent incisors	Estimated age range	
	Sheep	Goat
0 pair	Less than one years	Under 1years
1 pair	1-1½ years	1-2 years
2 pair	1½-2years	2-3 years
3 pair	2½-3years	3-4 years
4 pair	More than 3 years	More than four years
Broken	Aged	Aged

Source :( vatta *et al.*, 2006)

### Annex 3: Tick identification procedure

1. Ticks were removed from host skin carefully, not to damage the mouth parts, horizontally to the animal.

2. Code is given for ticks collected from each body regions with respect to date, site of attachments, their respective sex, age and breed of the animal.
3. The sample was put in universal bottle containing 70% ethanol and then it was transported to regional veterinary laboratory for identification purpose.
4. Ticks were identified to the genera and species level using stereomicroscope based on tick identification keys of (Kaiser, 1987 and Methyscae 1987).
5. Ticks were usually identified by the shape and length of capitulum, the colour of the body, the shape and markings on the scutum, host preferences and location on the host. Male and unengorged female ticks were easier to identify than engorged female ticks (Hendrix, 1998).

#### **Annex : 4 Materials General Equipment**

- ✓ White coveralls
- ✓ Boots,
- ✓ Datasheet
- ✓ Bag transporting tick samples
- ✓ Pencils and permanent marker pens for collection vials
- ✓ Thermometer (preferably digital)
- ✓ Mobile phone (optional)
- ✓ Forceps
- ✓ Magnifying glass to see ticks easier (optional)
- ✓ Garbage bags (to contain used drag cloths, coveralls, etc.

#### **DECLARATION**

I, the under signed, declare that the information presented here in my thesis is my original work, has not been published and not under consideration of publication in any journals and that all sources of materials used for this research have been duly acknowledged

Name: Mujahid Jewaro and Kufa Mustefa